JOURNAL OF CHROMATOGRAPHY B

VOL. 700, NOS. 1 + 2 24 OCTOBER 1997

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

Regular Papers

Purification and gas chromatographic–mass spectrometric characterization of non-methylene interrupted fatty acid incorporated in rat liver
by T. Tanaka, K. Shibata, H. Hino, T. Murashita, M. Kayama and K. Satouchi (Fukuyama, Japan) (Received 5 June 1997) . . . 1

Quantitative analysis of exogenous peptides in plasma using immobilized enzyme cleavage and gas chromatography–mass spectrometry with negative ion chemical ionization
by C.D. Márquez and M.-L. Lee (Austin, TX, USA), S.T. Weintraub (San Antonio, TX, USA) and P.C. Smith (Chapel Hill, NC, USA) (Received 14 May 1997) . . . 9

Simultaneous determination of polyamines, *N*-acetylated polyamines and the polyamine analogues BE-3-3-3 and BE-4-4-4-4 by capillary gas chromatography with nitrogen–phosphorus detection
by B. Dorhout, A.W. Kingma, E. de Hoog and F.A.J. Muskiet (Groningen, Netherlands) (Received 27 April 1997) . . . 23

Determination of endogenous levels of 13-*cis*-retinoic acid (isotretinoin), all-*trans*-retinoic acid (tretinoin) and their 4-oxo metabolites in human and animal plasma by high-performance liquid chromatography with automated column switching and ultraviolet detection
by R. Wyss and F. Bucheli (Basel, Switzerland) (Received 15 May 1997) . . . 31

Combination of high-performance liquid chromatography and thin-layer chromatography separation of five adducted nucleotides isolated from liver resulting from intraperitoneal administration with 7H-dibenzo[*c*,*g*]carbazole to mice
by P. O'Connor, S. Fremont, J. Schneider, H. Dowty, M. Jaeger, G. Talaska and D. Warshawsky (Cincinnati, OH, USA) (Received 22 May 1997) . . . 49

Bile acid kinetics in man studied by radio thin-layer chromatography and densitometry coupling
by D. Tonelli, E. Gattavecchia, G. Mazzella and A. Roda (Bologna, Italy) (Received 20 May 1997) . . . 59

Capillary electrophoresis of glutamate and aspartate in rat brain dialysate. Improvements in detection and analysis time using cyclodextrins
by W.H. Church (Harford, CT, USA), C.S. Lee (Greenville, NC, USA) and K.M. Dranchak (Harford, CT, USA) (Received 21 May 1997) . . . 67

Simultaneous determination of amphetamine and its analogs in human whole blood by gas chromatography–mass spectrometry
by P. Marquet, E. Lacassie, C. Battu, H. Faubert and G. Lachâtre (Limoges, France) (Received 13 May 1997) . . . 77

Hair analysis for drugs of abuse XIX. Determination of ephedrine and its homologs in rat hair and human hair
by Y. Nakahara and R. Kikura (Tokyo, Japan) (Received 12 June 1997) . . . 83

Determination of low levels of poly(ethylene glycol) 400 in plasma and urine by capillary gas chromatography–selected ion-monitoring mass spectrometry after solid-phase extraction
by C. Fakt and M. Ervik (Mölndal, Sweden) (Received 26 May 1997) . . . 93

High-performance liquid chromatography of unmodifed rosin and its applications in contact dermatology
by S. Sadhra (Birmingham, UK), C.N. Gray (Geelong, Australia) and I.S. Foulds (Birmingham, UK) (Received 2 May 1997) . 101

(Continued overleaf)

Contents (continued)

Analysis of a novel antiinflammatory agent, 1-(7-*tert.*-butyl-2,3-dihydro-3,3-dimethylbenzo[*b*]furan-5-yl)-4-cyclopropylbutan-1-one (PGV-20229), in plasma matrices by stable-isotope-dilution gas chromatography–mass spectrometry
by P.M. Knight, M.W. Scherz, C.A. Cruze and K.R. Wehmeyer (Mason, OH, USA) (Received 21 May 1997) 111

Screening for forensically relevant benzodiazepines in human hair by gas chromatography–negative ion chemical ionization-mass spectrometry
by V. Cirimele, P. Kintz and B. Ludes (Strasbourg, France) (Received 4 June 1997) 119

Validated assay for the quantification of anastrozole in human plasma by capillary gas chromatography–^{63}Ni electron capture detection
by M.J.H. Bock, I. Bara, N. LeDonne, A. Martz and M. Dyroff (Wilmington, DE, USA) (Received 19 February 1997) 131

Determination of the substance P receptor antagonist CP-122,721 in plasma by narrow-bore high-performance liquid chromatography–ionspray tandem mass spectrometry
by A. Kamel and C. Prakash (Groton, CT, USA) (Received 30 May 1997) 139

Simultaneous quantification of cefotaxime, desacetylcefotaxime, ofloxacine and ciprofloxacine in ocular aqueous humor and in plasma by high-performance liquid chromatography
by H.-J. Kraemer, R. Gehrke, A. Breithaupt and H. Breithaupt (Giessen, Germany) (Received 27 May 1997) 147

Enantioselective high-performance liquid chromatography determination of methadone enantiomers and its major metabolite in human biological fluids using a new derivatized cyclodextrin-bonded phase
by C. Pham-Huy, N. Chikhi-Chorfi, H. Galons, N. Sadeg, X. Laqueille, N. Aymard, F. Massicot, J.-M. Warnet and J.-R. Claude (Paris, France) (Received 23 May 1997) 155

Stereoselective determination of *R*-(+)- and *S*-(−)-remoxipride, a dopamine D_2-receptor antagonist, in human plasma by chiral high-performance liquid chromatography
by M.E. DePuy (West Point, PA, USA), J.L. Demetriades (Wayne, PA, USA) and D.G. Musson and J.D. Rogers (West Point, PA, USA) (Received 18 February 1997) 165

Quantitation of trimipramine enantiomers in human serum by enantioselective high-performance liquid chromatography and mixed-mode disc solid-phase extraction
by J. Liu and J.T. Stewart (Athens, GA, USA) (Received 23 May 1997) 175

Sensitive and rapid method for the simultaneous quantification of five antidepressants with their respective metabolites in plasma using high-performance liquid chromatography with diode-array detection
by G. Aymard, P. Livi, Y.T. Pham and B. Diquet (Paris, France) (Received 16 May 1997) 183

Determination of SDZ ICM 567 in blood and muscle microdialysis samples by microbore liquid chromatography with ultraviolet and fluorescence detection
by B. Ofner (Graz, Austria), A. Boukhabza, W. Pacha and Ch.V. Amsterdam (Basel, Switzerland) and R. Wintersteiger (Graz, Austria) (Received 2 April 1997) 191

Determination of propiomazine in rat plasma by direct injection on coupled liquid chromatography columns with electrochemical detection
by K. Benkestock (Stockholm, Sweden) (Received 27 May 1997) 201

Determination of 6,4′-bis-(2-imidazolinylhydrazone)-2-phenylimidazo[1,2-*a*]pyridine in plasma and whole blood by high-performance liquid chromatography
by T.S. Selvamuniandy, N. Mohamed, S.M. Mansor, N.K. Nair and V. Navaratnam (Penang, Malaysia) (Received 1 April 1997) 209

Stereospecific determination of mefloquine in biological fluids by high-performance liquid chromatography
by E. Souri and H. Farsam (Tehran, Iran) and F. Jamali (Edmonton, Canada) (Received 15 May 1997) 215

Confirmation of malachite green, gentian violet and their leuco analogs in catfish and trout tissue by high-performance liquid chromatography utilizing electrochemistry with ultraviolet-visible diode array detection and fluorescence detection
by L.G. Rushing and E.B. Hansen Jr. (Jefferson, AR, USA) (Received 27 May 1997) 223

Quantitative comparison on the refinement of horse antivenom by salt fractionation and ion-exchange chromatography
by T. Saetang, N. Treamwattana, P. Suttijitpaisal and K. Ratanabanangkoon (Bangkok, Thailand) (Received 4 March 1997) . . 233

Characterisation of a chromatographically produced anti-D immunoglobulin product
by M. Stucki, R. Moudry, C. Kempf, A. Omar, A. Schlegel and P.G. Lerch (Bern, Switzerland) (Received 27 May 1997) . . . 241

Analysis of the antidiabetic drug acarbose by capillary electrophoresis
by I. Rethfeld and G. Blaschke (Münster, Germany) (Received 5 June 1997) 249

Short Communications

Detection of microbial-derived fatty acids in carious dentin by gas chromatography and gas chromatography–mass spectrometry
by I. Brondz, H. Nordbø and A. Aamdal Scheie (Oslo, Norway) (Received 3 June 1997) 255

Measurement of oxalate in human plasma ultrafiltrate by ion chromatography
by M. Manoharan and P.O. Schwille (Erlangen, Germany) (Received 26 May 1997) 261

Microchromatographic quantitation of hemoglobin A levels in phenotypes of sickle cell-beta$^+$ thalassemia
by I. Sweeting, B.E. Serjeant, P.W. Thomas and G.R. Serjeant (Kingston, Jamaica) (Received 16 April 1997) 269

Thiopurine methyltransferase activity: new high-performance liquid chromatographic assay conditions
by Y. Medard, S. Nafa and E. Jacqz-Aigrain (Paris, France) (Received 29 April 1997) 275

Simultaneous determination of L-dopa and 3-O-methyldopa in human platelets and plasma using high-performance liquid chromatography with electrochemical detection
by F. Blandini, E. Martignoni, C. Pacchetti, S. Desideri, D. Rivellini and G. Nappi (Pavia, Italy) (Received 13 May 1997) . . . 278

High-performance thin-layer chromatographic determination of 5-methoxypsoralen in serum from patients
by B. Mignot, Y. Guillaume, S. Makki, E. Murret, E. Cavalli, T.T. Truong, M. Thomassin and C. Guinchard (Besançon, France) (Received 21 February 1997) 283

Erratum 286

Related Articles J. Chromatography A. Vols. 776–780 287

Author Index Volume 700 291

Compound Index Volume 700 295

The table of contents of *Journal of Chromatography* is included in ESTOC – Elsevier Science Tables of Contents service – which can be accessed on the World Wide Web at the following URL addresses:
http://www.elsevier.nl/locate/estoc or http://www.elsevier.com/locate/estoc